

KERNELS OF NUMERICAL PUSHFORWARDS

MIHAI FULGER AND BRIAN LEHMANN

ABSTRACT. Let $\pi : X \rightarrow Y$ be a morphism of projective varieties and consider the pushforward map $\pi_* : N_k(X) \rightarrow N_k(Y)$ of numerical cycle classes. We show that when the Chow groups of points of the fibers are as simple as they can be, then the kernel of π_* is spanned by k -cycles contracted by π .

1. INTRODUCTION

Let $\pi : X \rightarrow Y$ be a morphism of projective varieties over an algebraically closed field. The pushforward of cycles induces a map $\pi_* : N_k(X) \rightarrow N_k(Y)$ on numerical groups with $\mathbb{R}$ -coefficients. One would like to understand how $\ker(\pi_*)$ reflects the geometry of the map π . In the special case when $\alpha \in N_k(X)$ is the class of a closed subvariety Z , then α lies in the kernel of π_* precisely when $\dim \pi(Z) < \dim(Z)$. A similar statement holds when α is the class of an effective cycle. However, the geometry of arbitrary elements of $\ker(\pi_*)$ is more subtle, e.g. Example 1.1. We are interested in identifying classes of morphisms π for which the kernel of π_* reflects the “geometry” of the morphism in the simplest way: it is spanned by effective cycles contracted by π .

Perhaps the easiest example where this property fails is the following.

Example 1.1. Let E be an elliptic curve without complex multiplication and define $X = E \times E$. Then the Neron-Severi space $N_1(X)$ is 3-dimensional with a basis given by the classes F_1, F_2 of fibers of the two projections and by the diagonal class Δ . The pseudo-effective cone $\overline{\text{Eff}}_1(X)$ is a round cone.

Consider the first projection $\pi : X \rightarrow E$. The curves contracted by π are all numerically equivalent with class F_1 . However, the kernel of π is two-dimensional: it is generated by F_1 and by $\Delta - F_2$. In particular, the kernel is *not* spanned by the subvarieties contracted by π .

We expect that the failure in the previous example is caused by the fibers having large Chow groups. Our main result supports this.

Theorem 1.2. *Let $\pi : X \rightarrow Y$ be a dominant morphism of projective varieties over an uncountable algebraically closed field, with Y smooth. Suppose that every fiber F (over a closed point) of π satisfies $\dim_{\mathbb{Q}}(A_0(F)_{\mathbb{Q}}) = 1$. Then the kernel of π is spanned by effective π -contracted cycles.*

The corresponding statement for rational equivalence is well-established (see [BS83]). The main point is to translate between rational and numerical equivalence for the map π . The hypothesis is satisfied by any rationally chain connected variety. More generally, the varieties satisfying the hypothesis are regulated by Bloch’s Conjecture; for example, for a smooth surface S over $\mathbb{C}$ the condition $\dim_{\mathbb{Q}}(A_0(S)_{\mathbb{Q}}) = 1$ is expected to be equivalent to $\dim H^0(S, K_S) = 0$.

A few particular cases are worth special note:

Proposition 1.3 (cf. 3.8). *Let $\pi : X \rightarrow Y$ be a birational morphism over $\mathbb{C}$ with Y smooth. Then the kernel of π is spanned by effective π -contracted cycles.*

Corollary 1.4 (cf. 4.3). *Let $\pi : X \rightarrow Y$ be a morphism of projective varieties over $\mathbb{C}$ with Y smooth and with rationally chain connected general fiber. Then the kernel of π is spanned by effective π -contracted cycles.*

The second author is supported by NSF Award 1004363.

A related result for curve classes appears in [KM92, Corollary 12.1.5.1].

In a different direction, [DJV13] investigates (for an arbitrary morphism π) which π_* -contracted classes are actually linear combinations of classes of π -contracted subvarieties. The main conjecture of [DJV13] is that any π_* -contracted *pseudo-effective* class satisfies this linear combination property. In a forthcoming paper we use constructions from [FL14] to prove new cases of their conjecture.

2. BACKGROUND AND CONVENTIONS

By *variety* we mean a reduced, irreducible, separated scheme of finite type over an algebraically closed field of arbitrary characteristic. Unless otherwise stated, $\pi : X \rightarrow Y$ denotes a morphism of projective varieties. We say that a k -cycle Z is a $\mathbb{Z}$, $\mathbb{Q}$, or $\mathbb{R}$ - k -cycle depending on whether the coefficients lie in $\mathbb{Z}$, $\mathbb{Q}$, or $\mathbb{R}$. A cycle that is a positive combination of subvarieties is called *effective*. The group of all k - $\mathbb{Z}$ -cycles is denoted by $Z_k(X)$ and the Chow group of k - $\mathbb{Z}$ -cycles up to rational equivalence ([Ful84, §1]) is denoted by $A_k(X)$. A closed subscheme $Y \subset X$ determines a *fundamental $\mathbb{Z}$ -cycle* $[Y]$ (cf. [Ful84, §1.5]).

Let $\pi : X \rightarrow Y$ be a morphism of projective varieties and let Z be a k -cycle on X . We say that Z is π -*contracted* if for every component Z_i of Z we have $\dim(\pi(Z_i)) < \dim(Z_i)$.

We let $N_k(X)_{\mathbb{Z}}$ denote the quotient of $Z_k(X)$ by the relation of *numerical equivalence* as in [Ful84, Chapter 19]. $N_k(X)_{\mathbb{Z}}$ is a lattice inside the *numerical space* $N_k(X) := N_k(X)_{\mathbb{Z}} \otimes_{\mathbb{Z}} \mathbb{R}$. If Z is a k -cycle with $\mathbb{R}$ -coefficients, its class in $N_k(X)$ is denoted $[Z]$. The real Néron-Severi space of numerical classes of Cartier divisors is denoted $N^1(X)$. Its elements act as first Chern classes on the numerical groups $N_k(X)$.

For the rest of the paper, the term cycle will always refer to a cycle with $\mathbb{Z}$ -coefficients, and a numerical class will always refer to a class with $\mathbb{R}$ -coefficients, unless otherwise qualified.

3. THE GK PROPERTY

Definition 3.1. A morphism of projective varieties $\pi : X \rightarrow Y$ satisfies the *Geometric Kernel* (or GK) property if every class $\alpha \in N_k(X)$ with $\pi_*\alpha = 0$ lies in the vector space generated by the k -dimensional subvarieties which are contracted by π .

We begin with a few easy observations and important particular cases. Among them we will see that the GK property holds over $\mathbb{C}$ for birational maps π with a smooth base.

3.1. Compositions of morphisms. Note that if $\pi : X \rightarrow Y$ is a dominant morphism of projective varieties then $\pi_* : N_k(X) \rightarrow N_k(Y)$ is surjective, since any k -dimensional subvariety of Y is mapped onto by a k -dimensional subvariety of X .

Lemma 3.2. *The (k -th) GK property is stable under composition of dominant projective morphisms.*

Proof. Let $\pi_2 : X_2 \rightarrow X_1$ and $\pi_1 : X_1 \rightarrow X_0$ be dominant morphisms of projective varieties satisfying the GK property. For $i \in \{1, 2\}$, choose a finite set Σ_i of k -dimensional subvarieties of X_i whose classes generate $\ker \pi_{i*}$. Let Σ'_1 denote a set of k -dimensional subvarieties of X_2 such that each subvariety in Σ_1 is dominated under π_2 by a subvariety in Σ'_1 . Then the elements of $\Sigma'_1 \cup \Sigma_2$ are contracted by $\pi_1 \circ \pi_2$, and their classes span $\ker(\pi_1 \circ \pi_2)_*$. $\square$

Lemma 3.3. *If $\pi_1 : X_1 \rightarrow X_0$ and $\pi_2 : X_2 \rightarrow X_1$ are dominant morphisms such that $\pi_1 \circ \pi_2$ satisfies the GK property, then π_1 satisfies the GK property.*

Proof. There is a surjection $\pi_{2*} : \ker(\pi_{1*} \circ \pi_{2*}) \rightarrow \ker(\pi_{1*})$, and the desired conclusion follows. $\square$

3.2. The GK property for projective bundles and birational maps.

Proposition 3.4. *Projective bundle maps satisfy the GK property.*

Proof. Let $X = \mathbb{P}_Y(E)$ for some vector bundle E of rank r on Y , and denote by $\pi : X \rightarrow Y$ the bundle map. Let $\xi = [c_1(\mathcal{O}_{\mathbb{P}(E)}(1))] \in N^1(X)$. Up to twisting E , we can assume that ξ is ample.

Suppose $\pi_*\alpha = 0$ for some $\alpha \in N_k(X)$. Using [Ful84, Theorem 3.3.(b)], write

$$\alpha = \sum_{i=0}^{r-1} \xi^{r-1-i} \cdot \pi^* a_{k-i}$$

for some classes $a_{k-i} \in N_{k-i}(Y)$. Since $\pi_*\xi^{r-1} = [Y] \in N_{\dim Y}(Y)$, the condition $\pi_*\alpha = 0$ is equivalent to $a_k = 0$. Thus, it suffices to see that for each $i > 0$ the term $\xi^{r-1-i} \cdot \pi^* a_{k-i}$ is represented by a linear combination of cycles contracted by π .

Let m be a sufficiently positive integer so that $m\xi$ is very ample. For any cycle Z , the numerical class $\xi \cdot [Z]$ is represented by $\frac{1}{m}H|_Z$ where H is a general element of $|m\xi|$. Using this construction repeatedly, we see that $\xi^{r-1-i} \cdot \pi^* a_{k-i}$ is represented by a cycle with positive relative dimension over Y for any $i > 0$ and any class $a \in N_{k-i}(Y)$. $\square$

Remark 3.5. Essentially the same argument shows that Grassmann bundle maps satisfy the GK property.

Lemma 3.6. *Let $\pi : X \rightarrow Y$ be the blow-up of a smooth variety Y along a smooth center. Then π satisfies the GK property.*

Proof. Let Z be the blow-up center so that we have a cartesian diagram:

$$\begin{array}{ccc} \tilde{Z} & \xrightarrow{j} & X \\ f \downarrow & & \downarrow \pi \\ Z & \xrightarrow{i} & Y \end{array}$$

with $f : \tilde{Z} \rightarrow Z$ the projective bundle $\mathbb{P}(N_Z^\vee Y)$. By [Ful84, Proposition 6.7], any cycle α on X is of the form $j_*\tilde{z} + \pi^*y$. The condition $\pi_*\alpha = 0$ implies $y = -i_*f_*\tilde{z}$. By [Ful84, Proposition 6.7.(a) and Example 3.3.3], we have $\alpha = j_*\beta$ with $f_*\beta = 0$. Proposition 3.4 completes the proof. $\square$

Remark 3.7. The pullback map π^* appearing in the proof of Proposition 3.4 is well-defined for Chow groups and respects numerical equivalence because π is smooth (cf. [Ful84, Example 19.2.3]). This is of slightly different flavor from the pullback π^* used in the proof of Lemma 3.6. Here, π^* is well-defined on Chow groups and respects numerical equivalence because Y is smooth (cf. [Ful84, Example 19.1.6]).

We caution the reader that it is not clear that the flat (non-smooth) pullback for Chow groups from [Ful84, §1.7] respects numerical equivalence. We will not be concerned with this difficulty here.

Proposition 3.8. *Suppose that $\pi : X \rightarrow Y$ is a birational morphism of varieties over $\mathbb{C}$ with Y smooth. Then π satisfies the GK property.*

Note that the case of divisors is well-known; in fact, the Negativity of Contraction lemma proves the GK property for divisor classes when we have a birational map in arbitrary characteristic with a $\mathbb{Q}$ -factorial base.

Proof. The birational map π is dominated by a composition of blow-ups of smooth varieties along smooth centers. The result follows by Lemma 3.2, Lemma 3.3 and Lemma 3.6. $\square$

Example 3.9. The smoothness of Y in the proposition is necessary. For example if Z is the projective cone over $E \times E$, where E is an elliptic curve, and $\tau : X \rightarrow Z$ is the blow-up of the vertex

with exceptional divisor $D \simeq E \times E$, then X has a structure of a projective bundle over $E \times E$ with bundle map f . Let $\pi : X \rightarrow Y$ be the contraction of a fiber of the first projection $D = E \times E \rightarrow E$. Then

$$\pi_*(D \cdot f^*(\Delta - F_2)) = 0,$$

where Δ is the class of the diagonal on $E \times E$ and F_2 is a fiber of the second projection. However the only effective curve contracted by π is $D \cdot f^*F_1$, and $D \cdot f^*(\Delta - F_2)$ is not a multiple of it. $\square$

3.3. The GK property and Chow groups of points. We first look to understand the relative behavior of Chow groups under a morphism π . For this we use a variant of the Bloch–Srinivas theorem presented in [Voi07, Theorem 10.19].

Theorem 3.10. *Let $\pi : X \rightarrow Y$ be a projective map of quasi-projective schemes of finite type over an uncountable algebraically closed field with Y integral. Let Z be a k -cycle on X with rational equivalence class τ . Suppose there is a subscheme X' of X such that for a general smooth closed point $y \in Y$, the restriction $\tau|_{X_y}$ vanishes in $A_*(X_y - X'_y)_{\mathbb{Q}}$.*

Then there is a positive integer m , a cycle Z_1 in X' , and a cycle Z_2 whose support does not dominate Y such that $mZ \sim Z_1 + Z_2$ in $A_k(X)$.

Since this statement is undoubtedly well-known to experts, we only give a brief verification.

Proof. Let $\overline{X}$ be a projective closure of X , and let H be a very ample divisor on $\overline{X}$. By $|rH|$ we denote the complete linear series on $\overline{X}$. If W is a projective subvariety of X , then $|rH|_W$ naturally surjects onto $|rH|_W$ for all $r \gg 0$.

Most of the proof of [Voi07, Theorem 10.19], whose notation we retain, goes through. We list the changes:

- The condition $\tau|_{X_y} = 0 \in A_*(X_y - X'_y)_{\mathbb{Q}}$ allows torsion $\mathbb{Z}$ -classes. Consequently the relative Hilbert schemes H_i used in the proof of [Voi07] are now defined to parameterize rational equivalence relations between sZ_y and cycles in X'_y for all positive integers s .
- The resolution of the subvariety H mapping generically finitely onto Y is replaced by a nonsingular alteration ([dJ96]).
- The divisors $\text{div}(\phi_{y,l})$ on $W_{y,l}$ are differences of elements of $|rH|_{W_{y,l}}$, hence they can be seen as differences of restrictions of elements of $|rH|$ and parameterized accordingly. Then the divisors $\mathcal{D}_l$ can be constructed directly as differences of restrictions to $\mathcal{W}_l$ of divisors on $\mathbb{P}(|rH|) \times H$ for some $r \gg 0$, in particular they are Cartier.
- As in the proof of [Voi07, Lemma 10.22], one uses Grauert's theorem to show that the Cartier divisors $\mathcal{D}_l$ are pullbacks from Y , at least over an open subset of Y .

$\square$

We then immediately obtain:

Corollary 3.11. *Let $\pi : X \rightarrow Y$ be a projective map of quasi-projective schemes of finite type over an uncountable algebraically closed field, with Y integral of dimension k . Suppose that for a fiber X_y over a general closed point y we have $\dim_{\mathbb{Q}}(A_0(F)_{\mathbb{Q}}) = 1$. Then if Z is a k -cycle on X such that $\pi_*Z \sim_{\mathbb{Q}} 0$, there is a positive integer m and a π -contracted cycle Z' such that $mZ \sim_{\mathbb{Z}} Z'$.*

Proof. Using the restriction exact sequence for Chow groups, we may shrink Y to assume that it is smooth.

If every component of Z is contracted by π , set $Z' = Z$. Otherwise, Z must have some component Z_1 that surjects onto Y . Since a general fiber X_y has $\text{rank}(A_0(X_y)) = 1$, the restriction of Z to $A_0(X_y - Z_{1,y})_{\mathbb{Q}}$ must vanish. Thus, up to adding some π -contracted cycle, a multiple of Z is rationally equivalent to a multiple of Z_1 . But of course this multiple must be 0 since it pushes forward to 0. $\square$

Remark 3.12. Consider a dominant morphism of projective varieties $\pi : X \rightarrow Y$ with Y smooth. Then there is an induced surjective morphism

$$A_k(X)_{\mathbb{Q}} \cap \ker \pi_*^{\text{Chow}_{\mathbb{Q}}} \rightarrow N_k(X)_{\mathbb{Q}} \cap \ker \pi_*^{\text{num}_{\mathbb{Q}}}.$$

(If π has relative dimension e and if h is an ample class on X normalized so that $\pi_*(h^e \cap [X]) = [Y]$, then $\pi_*(h^e \cdot \pi^* \alpha) = \alpha$ for any $\alpha \in A_k(Y)$. This and [Ful84, Example 19.1.6] imply that π_* maps numerically trivial Chow classes on X surjectively onto numerically trivial Chow classes on Y . Conclude by the Snake Lemma.)

Lemma 3.13. *Let $\pi : X \rightarrow Y$ be a surjective map of projective varieties with Y smooth, and let Z be a k - $\mathbb{R}$ -cycle on X such that $\pi_*[Z] = 0 \in N_k(Y)$. Then there is some k - $\mathbb{R}$ -cycle $Z' \equiv Z$ such that $\pi_*Z' = 0$ as cycles.*

Proof. Since π_* is defined over $\mathbb{Z}$, we can replace $\mathbb{R}$ with $\mathbb{Q}$ throughout. By Remark 3.12, there exists a cycle $Z_1 \equiv Z$ such that $\pi_*Z_1 \in \text{Rat}_k(Y)$, where $\text{Rat}_k(Y)$ denotes the group of k - $\mathbb{Q}$ -cycles rationally equivalent to zero on Y . It is then enough to show that π_* maps $\text{Rat}_k(X)$ onto $\text{Rat}_k(Y)$. By [Ful84, §1.3], if $A \in \text{Rat}_k(Y)$, then there exist finitely many $k+1$ -dimensional subvarieties V_i of Y and elements $r_i \in R(V_i)^*$ such that $A = \sum_i \text{div}(r_i)$. Let W_i be $k+1$ -dimensional subvarieties of X mapping generically finitely onto V_i , and denote $d_i = \deg(W_i/V_i)$. Then $A' = \sum_i \frac{1}{d_i} \text{div}(r_i) \in \text{Rat}_k X$ satisfies $\pi_*A' = A$ by [Ful84, Proposition 1.4.(b)]. $\square$

4. THE MAIN RESULT

Proof of Theorem 1.2. It suffices to consider the case when $k \leq \dim Y$. Let Z be any k -cycle such that $\pi_*[Z] = 0$. Apply Lemma 3.13 to replace Z by the numerically equivalent cycle Z' as in the statement. Thus Z' is a sum of cycles Z_i such that each Z_i satisfies $\pi_*Z_i = 0$ and either:

- $|Z_i|$ is π -contracted, or
- $\pi(|Z_i|)$ is an irreducible k -dimensional subvariety of Y .

Let $p_i : W_i \rightarrow T_i$ be the base change of π to $\pi(Z_i)$. Applying Corollary 3.11 to p_i , we see that each Z_i is $\mathbb{Q}$ -rationally equivalent to a π -contracted cycle (as cycles on W_i , and hence after pushforward as cycles on X). Thus Z is numerically equivalent to a sum of π -contracted cycles. $\square$

Remark 4.1. In a slightly different setting, [KM92, Corollary 12.1.5.1] shows that if $\pi : X \rightarrow Y$ is a projective morphism with connected fibers between algebraic spaces with rational singularities, and $R^1\pi_*\mathcal{O}_X = 0$, then the GK property holds for π when $k = 1$.

Remark 4.2. The condition $\dim_{\mathbb{Q}}(A_0(F)_{\mathbb{Q}}) = 1$ can be relaxed, asking instead that the closure of the locus in Y where it fails have dimension at most $k - 1$.

Corollary 4.3. *Suppose that $\pi : X \rightarrow Y$ is a morphism of projective varieties over $\mathbb{C}$ such that Y is smooth and the general fiber of π is rationally chain connected. Then π satisfies the GK property.*

Proof. By Proposition 3.8 and Lemmas 3.2 and 3.3, we may replace π by a birationally equivalent morphism. Since π is flat over an open subset, there is a rational map $Y \dashrightarrow \text{Hilb}(X)$ induced by the fibers. Since we may replace Y by a birational model Y' resolving this map (cf. Proposition 3.8) and may replace X by the main component of $X \times_Y Y'$ for X , we may assume that π is flat (and Y is still smooth). Note that the general fiber is unchanged by the flattening operation. By Theorem 1.2 it suffices to show that rational chain connectedness is a flat deformation invariant. It suffices to consider the case when we have a flat morphism $\pi : X \rightarrow C$ for a curve C where the general fiber is RCC. Let F_0 denote the special fiber; we must show any two points x, x' of F_0 are connected by a chain of rational curves. Choose a curve C' on X that dominates C and contains x and x' . For a general fiber F , there is a rational chain in F connecting all the points of $C' \cap F$. Using the uncountability of $\mathbb{C}$, one sees that such rational chains form a family of curves which

dominates the base. By taking a closure and applying [Kol96, II.2.4 Corollary], we obtain a chain of rational curves in F_0 which connects x and x' . $\square$

Example 4.4. Let S be a smooth surface such that $A_0(S) = \mathbb{Z}$. By the work of [Mum68] and [Roĭ72], this implies that $p_g = 0$ and $\text{Alb}(S)$ is trivial. Examples include any rational surface S and conjecturally any surface with $q = p_g = 0$.

Suppose that Y is another smooth surface. We claim that there is an isomorphism

$$N_2(S \times Y) \cong \mathbb{R} \oplus (N_1(S) \otimes N_1(Y)) \oplus \mathbb{R}.$$

For surfaces over $\mathbb{C}$, this follows easily from Hodge theory and the Kunneth formula. (In fact, this argument also works for any surface S satisfying $q = p_g = 0$.) We present an alternative “algebraic” approach valid over any uncountable algebraically closed field.

Let $\pi : S \times Y \rightarrow Y$ be the projection map. By Theorem 1.2 the kernel of $\pi_* : N_2(S \times Y) \rightarrow \mathbb{R}$ is spanned by π -contracted irreducible surfaces. First note that any irreducible surface contracted to a point by π is a fiber.

Second, suppose that the image of a π -contracted surface T is a curve C on Y . Let C' be a normalization of C . We show that $N_2(S \times C') \cong N_1(S) \times \mathbb{R}$. If an effective surface T' on $S \times C'$ does not dominate S , then it is the pullback of a divisor on S . If it does dominate S , then it induces a morphism $S \rightarrow \text{Jac}(C')$. But by assumption on the Albanese map this morphism is trivial. So after twisting by the pullback of a line bundle from S , the divisor T' is the pullback of a divisor on C' .

Putting these together, we see N_2 is spanned by three kinds of classes: fibers of π_1 , fibers of π_2 , and products $\pi_1^*D \cdot \pi_2^*E$ for a divisor D on S and E on Y . The natural map $\pi_1^*N_1(S) \otimes \pi_2^*N_1(Y) \rightarrow N_2(S \times Y)$ is easily seen to be injective. $\square$

REFERENCES

- [BS83] Spencer Bloch and Vasudevan Srinivas, *Remarks on correspondences and algebraic cycles*, Amer. J. Math. **105** (1983), no. 5, 1235–1253.
- [dJ96] Aise Johan de Jong, *Smoothness, semi-stability and alterations*, Inst. Hautes Études Sci. Publ. Math. (1996), no. 83, 51–93.
- [DJV13] Olivier Debarre, Zhi Jiang, and Claire Voisin, *Pseudo-effective classes and pushforwards*, Pure Appl. Math. Q. **9** (2013), no. 4, 643–664.
- [FL14] Mihai Fulger and Brian Lehmann, *Positive cones of dual cycle classes*, 2014, arXiv:1408.5154.
- [Ful84] William Fulton, *Intersection theory*, Ergebnisse der Mathematik und ihrer Grenzgebiete (3) [Results in Mathematics and Related Areas (3)], vol. 2, Springer-Verlag, Berlin, 1984.
- [KM92] János Kollár and Shigefumi Mori, *Classification of three-dimensional flips*, J. Amer. Math. Soc. **5** (1992), no. 3, 533–703.
- [Kol96] János Kollár, *Rational curves on algebraic varieties*, Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge. A Series of Modern Surveys in Mathematics [Results in Mathematics and Related Areas. 3rd Series. A Series of Modern Surveys in Mathematics], vol. 32, Springer-Verlag, Berlin, 1996.
- [Mum68] David Mumford, *Rational equivalence of 0-cycles on surfaces*, J. Math. Kyoto Univ. **9** (1968), 195–204.
- [Roĭ72] A. A. Roĭtman, *Rational equivalence of zero-dimensional cycles*, Mat. Sb. (N.S.) **89(131)** (1972), 569–585, 671.
- [Voi07] Claire Voisin, *Hodge theory and complex algebraic geometry. II*, Cambridge Studies in Advanced Mathematics, vol. 77, Cambridge University Press, Cambridge, 2007.

DEPARTMENT OF MATHEMATICS, PRINCETON UNIVERSITY, PRINCETON, NJ 08544

INSTITUTE OF MATHEMATICS OF THE ROMANIAN ACADEMY, P. O. BOX 1-764, RO-014700, BUCHAREST, ROMANIA

E-mail address: `afulger@princeton.edu`

DEPARTMENT OF MATHEMATICS, BOSTON COLLEGE, CHESTNUT HILL, MA 02467

E-mail address: `lehmannb@bc.edu`